On the 3x + 1 conjecture.

Peter Hellekalek*

July 26, 2016

Abstract

In this paper, we discuss the well known 3x + 1 conjecture in form of the accelerated Collatz function T defined on the positive odd integers. We present a sequence of quotient spaces and further, an invertible map, which are intrinsically related to the behavior of T. This approach allows to express the 3x + 1 conjecture in form of equivalent problems, which might be more accessible than the original conjecture.

1 Introduction

Let \mathbb{N}_0 stand for the nonnegative integers, $\mathbb{N}_0 = \{0, 1, \ldots\}$, let $\mathbb{U} = 2\mathbb{N}_0 + 1$ be the set of odd positive integers, and let

$$T: \mathbb{U} \to \mathbb{U}, \quad Tx = (3x+1)2^{-\nu_2(3x+1)},$$

where $\nu_2(y)$ denotes the exponent of the largest power of 2 that divides the integer y. The map T is called the reduced or accelerated Collatz function in the literature (see [1]). Hence, T17 = 13, T13 = 5, and T5 = 1.

The 3x+1 conjecture, also known as the Collatz conjecture, states that, starting from any $x \in \mathbb{U}$, by iterating T we will eventually end up in the number 1. In other words, for every $x \in \mathbb{U}$, there exists $k = k(x) \in \mathbb{N}_0$ such that $T^k x = 1$. Here, T^0 stands for the identity map, and, for $k \in \mathbb{N}$, T^k is defined recursively by $T^k = T \circ T^{k-1}$. To give an example, $T^3 17 = 1$. We refer the reader to the comprehensive monograph [1] for details on the 3x+1 conjecture and for the numerous aspects that have been studied in this context.

^{*}The author is supported by the Austrian Science Fund (FWF): Project F5504-N26, which is a part of the Special Research Program "Quasi-Monte Carlo Methods: Theory and Applications

²⁰¹⁰ Mathematics Subject Classification: Primary 11T71; Secondary 94A60. Key words and phrases: 3x+1, Collatz problem, Syracuse problem

In this paper, we show how to associate with T an invertible map T^* on a certain quotient space that consists of equivalence classes of odd integers. The properties of T^* reflect the behavior of T, see Theorem 2.16. Further, we exhibit several statements that are equivalent to the 3x+1 conjecture, see Corollary 2.13. In the appendix, we present additional concepts to describe the action of T on \mathbb{U}_0 .

Our approach for the accelerated Collatz map T may be of interest for any such many-to-one map with a unique fixed point.

2 Results

For the sake of better readability, we will write Tx instead of T(x) for the image of x under T. The same slight abuse of notation will apply to the functions S and f below. All other functions will be written as usual.

Remark 2.1. The outline of our approach is the following:

- 1. First, we introduce a map $S: \mathbb{U} \to \mathbb{U}$ that allows to describe the inverse image $T^{-1}\{y\}$ of a point $y \in \mathbb{U}$, $y \not\equiv 0 \pmod{3}$, completely.
- 2. The next idea is to restrict T to the subset \mathbb{U}_0 of \mathbb{U} . T is surjective on \mathbb{U}_0 , hence we may introduce the *inverse map* τ , inverse in the sense $T \circ \tau$ being the identity map on \mathbb{U}_0 .
- 3. We then study a sequence of equivalence relations " \sim_n ", $n \geq 0$, which yields, for every $x \in \mathbb{U}_0$, an increasing sequence of equivalence classes $([x]_n)_{n\geq 0}$ and an associated decreasing sequence of positive integers $(\delta_n(x))_{n\geq 0}$.
- 4. The next idea is to study an equivalence relation " \sim_{∞} " on \mathbb{U}_0 , which leads to a partition of \mathbb{U}_0 into equivalence classes $[x]_{\infty}$, $x \in \mathbb{U}_0$, and to minimal elements $\delta_{\infty}(x)$ with the property $\delta_{\infty}(x) = \lim_{n \to \infty} \delta_n(x)$. Further, a bijective map T^* that mimics the behavior of T may be defined on the quotient space $\mathbb{U}_0/\sim_{\infty}$.
- 5. The 3x+1 conjecture is then equivalent to $\mathbb{U}_0 = [1]_{\infty}$, and also equivalent to $\delta_{\infty}(x) = 1$, for all $x \in \mathbb{U}_0$.
- 6. In the appendix, we analyze a closely related equivalence relation on \mathbb{U}_0 , which results in a partition of \mathbb{U}_0 into T-invariant subsets.
- 7. Also in the appendix, we extend the map T to an invertible map f on \mathbb{U}_0 . In addition, we provide some concepts for "bookkeeping" concerning the classes $[f^k x]_n$ and the positive integers $\delta_n(f^k x)$, where $k \in \mathbb{Z}$ and $n \geq 0$.

Remark 2.2. Our abstract approach sheds some light on the behavior of the map T. Why do we fail to prove the 3x + 1 conjecture or, at least, some partial results? What is missing in our study are quantitative results, for example

- 1. A description of the growth behavior of the classes $[1]_n$, in dependence of n. (This might yield a result on the density of $[1]_{\infty}$ in \mathbb{U}_0 .)
- 2. Number-theoretical arguments proving –for an appropriate notion of distance– that the distance between the class $[1]_{\infty}$ and each class $[x]_{\infty}$ can be made arbitrarily small. Equivalently, one could try to show that the assumption $\delta_{\infty}(x) > 1$ leads to a contradiction. (This would yield $[1]_{\infty} = [x]_{\infty}$ for all $x \in \mathbb{U}_0$, thereby proving the 3x+1 conjecture.)
- 3. (Number-theoretic) Arguments showing that there are no periodic points $[x]_{\infty}$ of T^* with a period larger or equal to 2. (This would imply that there is no periodic point of T with period larger or equal to 2, which is yet unknown.)

After this outline of concepts and shortcomings in our approach, let us look at the details.

 $Remark\ 2.3.$ The following properties of the map T are well known and elementary to prove.

1. From the definition of T, we derive the equivalence

$$y = Tx \Leftrightarrow 3x + 1 = y2^{\nu_2(3x+1)}.$$
 (1)

2. For every $y \in \mathbb{U}$ with $y \equiv 0 \pmod{3}$,

$$T^{-1}\{y\} = \emptyset.$$

This follows from (1) for the simple reason that $3x + 1 \equiv 1 \pmod{3}$, whereas $y2^{\nu_2(3x+1)} \equiv 0 \pmod{3}$.

3. The element 1 is the unique fixed point of T, i.e.

$$\{x \in \mathbb{U} : Tx = x\} = \{1\}.$$

Again, this is a direct consequence of (1).

The following map allows to describe the behaviour of T.

Definition 2.1. We define $S: \mathbb{U} \to \mathbb{U}$ as Sx = 4x + 1.

The map S permutes the residue classes modulo 3: if $x \equiv a \pmod{3}$, then $Sx \equiv a+1 \pmod{3}$. This simple property will prove to be essential for defining an inverse map associated with T, see Definition 2.3.

The next lemma is part of the 'folklore' in the 3x + 1 community. We present a simple proof, for the sake of completeness.

Lemma 2.1. For all $x \in \mathbb{U}$, we have

$$Tx = T(Sx). (2)$$

Proof. We have

$$T(Sx) = T(4x+1) = (3x+1)2^{2-\nu_2(12x+4)}$$
.

Trivially,
$$\nu_2(12x+4) = 2 + \nu_2(3x+1)$$
.

Corollary 2.2. Lemma 2.1 implies for all $x \in \mathbb{U}$ that x and its iterates $S^k x$ are mapped to Tx:

$$\forall x \in \mathbb{U}, \forall k \ge 0: \quad Tx = T(S^k x).$$

In other words, $T = T \circ S^k$ on \mathbb{U} , for all $k \geq 0$.

Remark 2.4. By induction for k we see that

$$\forall x \in \mathbb{U}, \forall k > 0: \quad S^k x = 4^k x + (4^k - 1)/3.$$

Definition 2.2. For $y \in \mathbb{U}$ with $y \not\equiv 0 \pmod{3}$, let $\xi(y)$ denote the smallest element of \mathbb{U} that is mapped to y by T:

$$\xi(y) = \min\{x \in \mathbb{U} : Tx = y\}.$$

Lemma 2.3. Let $y \in \mathbb{U}$ with $y \not\equiv 0 \pmod{3}$. Then $\xi(y)$ is given as follows.

- 1. If $y \equiv 1 \pmod{3}$, then $\xi(y) = (4y 1)/3$.
- 2. If $y \equiv 2 \pmod{3}$, then $\xi(y) = (2y 1)/3$.

Proof. Suppose that $y \equiv 1 \pmod{3}$. Then equivalence (1) implies that $\nu_2(3x+1)$ has to be even. The smallest solution in \mathbb{U} to (1) is the number x with the property $\nu_2(3x+1)=2$. This yields $\xi(y)=(4y-1)/3$.

The case $y \equiv 2 \pmod{3}$ is treated in the same manner.

Corollary 2.4. Suppose that $y \equiv 0 \pmod{3}$. Then, for $z \in \{Sy, S^2y\}$, the preimage $T^{-1}\{z\}$ is non-void.

Lemma 2.5. The set $T^{-1}\{Tx\}$ of those elements z of \mathbb{U} that are mapped to Tx is given by $\xi(Tx)$ and its iterates under S:

$$\{z \in \mathbb{U} : Tz = Tx\} = \{S^k \xi(Tx) : k \ge 0\}.$$

Proof. Suppose first that $y = Tx \equiv 1 \pmod{3}$, and assume that Tz = Tx. It follows from (1) that $\nu_2(3z+1) \in \{2,4,6,\ldots\}$. If $\nu_2(3z+1) = 2$, then from Lemma 2.3, Part 1, it follows that $z = \xi(Tx)$. If $\nu_2(3z+1) = 4$, then

$$3z + 1 = 2^4y = 2^2(3\xi(Tx) + 1).$$

This implies $z = S\xi(Tx)$.

In the general case, if $\nu_2(3z+1)=2+2k$, with $k\geq 1$, we have

$$3z + 1 = y2^{2+2k} = (3\xi(Tx) + 1)4^k.$$

It follows that

$$z = 4^k \xi(Tx) + (4^k - 1)/3,$$

from which we derive by Remark 2.4 that $z = S^k \xi(Tx)$.

Lemma 2.6. The set $\mathbb{U}_0 = \{x \in \mathbb{U} : x \not\equiv 0 \pmod{3}\}$ has the properties $T\mathbb{U} = \mathbb{U}_0$ and $T\mathbb{U}_0 = \mathbb{U}_0$. In particular, the map $T : \mathbb{U}_0 \to \mathbb{U}_0$ is surjective.

Proof. By Remark 2.3(2), we have $T\mathbb{U} \subseteq \mathbb{U}_0$, hence $T\mathbb{U}_0 \subseteq \mathbb{U}_0$. If $y \in \mathbb{U}_0$, then by Lemma 2.3 there exists $x \in \mathbb{U}$ such that Tx = y. Due to Lemma 2.1, we may assume $x \in \mathbb{U}_0$.

Corollary 2.7. In order to prove the 3x+1 conjecture, it suffices to restrict the map T to the set \mathbb{U}_0 .

Hence, from now on, we will study the 3x+1 conjecture for the surjective map $T: \mathbb{U}_0 \to \mathbb{U}_0$. We note that the surjectivity of T implies for all subsets B of \mathbb{U}_0 ,

$$T(T^{-1}B) = B. (3)$$

We employ the well-ordering principle to define some sort of inverse map associated with T.

Definition 2.3. For $x \in \mathbb{U}_0$, define the (quasi-)inverse function τ of T as follows:

$$\tau(x) = \min\{z \in \mathbb{U}_0 : Tz = x\}.$$

The reader should note that, for $\xi(x) \in \mathbb{U}_0$, $\tau(x) = \xi(x)$, whereas for $\xi(x) \equiv 0 \pmod{3}$, we have $\tau(x) = S\xi(x)$. To give an example, $\tau(5) = 13$, whereas $\xi(5) = 3$. Further, $T \circ \tau$ is the identity map on \mathbb{U}_0 , whereas, in general, $\tau(Tx) \neq x$.

The next idea is to generate a series of equivalence relations and, hence, a series of quotient spaces and of partitions of \mathbb{U}_0 .

Definition 2.4. For $x, y \in \mathbb{U}_0$ and $n \in \mathbb{N}_0$, we define the relation " \sim_n " on \mathbb{U}_0 as

$$x \sim_n y \Leftrightarrow T^n x = T^n y$$

Further, we put $[x]_n = \{z \in \mathbb{U}_0 : T^n z = T^n x\}$, and $\delta_n(x) = \min[x]_n$.

For all $x \in \mathbb{U}_0$, and all $n \geq 0$, we have $x \in [x]_n$, hence $[x]_n \neq \emptyset$ and $\delta_n(x) \leq x$. The set $[x]_0$ consists of the single point x.

Lemma 2.8. For all $x \in \mathbb{U}_0$ and for all $n \in \mathbb{N}_0$, the following holds.

- 1. The relation ' \sim_n ' is an equivalence relation on \mathbb{U}_0 and the set $[x]_n$ is the equivalence class of x with respect to this equivalence relation. Further, $[x]_n = [\delta_n(x)]_n$.
- 2. We have strict inclusion $[x]_n \subset [x]_{n+1}$.
- 3. For all $n \geq 1$ and all $k \in \mathbb{N}_0$ such that $S^k x \in \mathbb{U}_0$,

$$[x]_n = [S^k x]_n.$$

4. We have

$$1 \le \delta_n(x) \le \delta_{n-1}(x) \le \dots \le \delta_1(x) \le \delta_0(x) = x.$$

5. For all $x \in \mathbb{U}_0$ and all $n \geq 0$,

$$T^{-1}[Tx]_n = [x]_{n+1}.$$

Proof. Ad 1. This is easy to verify.

Ad 2. The inclusion $[x]_n \subseteq [x]_{n+1}$ is trivial. It follows from the definition of these two sets. In order to prove strict inclusion, put $y = T^n x$. If $y \equiv 1 \pmod{3}$, then let $z \in \mathbb{U}_0$ be such that $T^n z = Sy$. Hence, $z \notin [x]_n$. On the other hand, $T^{n+1}x = Ty = T(Sy) = T^{n+1}z$, which implies $z \in [x]_{n+1}$. If $y \equiv 2 \pmod{3}$, then let $z \in \mathbb{U}$ be such that $T^n z = S^2 y$. As above, we derive $z \in [x]_{n+1} \setminus [x]_n$.

Ad 3. From Corollary 2.2 it follows that, for all $n \geq 1$, we have the identity $T^n = T^n \circ S^k$ on \mathbb{U} and, hence, also on \mathbb{U}_0 . This implies $x \sim_n S^k x$, for all those $k \geq 0$ where $S^k x \in \mathbb{U}_0$.

Ad 4. Trivial.

Ad 5. Let $z \in T^{-1}[Tx]_n$. Then $Tz \in [Tx]_n$, which implies $T^n(Tz) = T^{n+1}z = T^n(Tx) = T^{n+1}x$. Hence, $z \in [x]_{n+1}$. This yields $T^{-1}[Tx]_n \subseteq [x]_{n+1}$. For the converse, if $z \in [x]_{n+1}$, then $T^{n+1}z = T^n(Tz) = T^{n+1}x = T^n(Tx)$, which implies $Tz \in [Tx]_n$. As a consequence, $z \in T^{-1}[Tx]_n$. We derive $[x]_{n+1} \subseteq T^{-1}[Tx]_n$.

Corollary 2.9. For all $x \in \mathbb{U}_0$, and for all $n \geq 0$,

$$T[x]_{n+1} = [Tx]_n, \quad T^{-1}[x]_n = [\tau(x)]_{n+1}, \quad T[\tau(x)]_{n+1} = [x]_n.$$

This is due to the surjectivity of T, see identity (3). In addition, $T[x]_0 = [Tx]_0$.

Corollary 2.10. For all $n \geq 0$, we may partition \mathbb{U}_0 as follows:

$$\mathbb{U}_0 = \bigcup_{x \in \mathbb{U}_0} [x]_n. \tag{4}$$

Due to the strict inclusion $[x]_n \subset [x]_{n+1}$, if we pass from n to n+1, this will result in a 'reduction' in the number of different equivalence classes. Hence, if n increases, we get less and less elements in the partitions (4) of \mathbb{U}_0 . As we will see in Corollary 2.13, the 3x+1 conjecture is equivalent to a collapse of this sequence of nested partitions to a trivial partition of \mathbb{U}_0 consisting of a single set.

Corollary 2.11. For all $x \in \mathbb{U}_0$, the limit $\lim_{n\to\infty} \delta_n(x)$ exists. This is due to the fact that the sequence of positive integers $(\delta_n(x))_{n\geq 0}$ is decreasing and bounded from below by 1, hence convergent.

We observe that the 3x+1 conjecture is equivalent to $\lim_{n\to\infty} \delta_n(x) = 1$ for all x in \mathbb{U}_0 . It is also equivalent to $1 \in \bigcup_{n>0} [x]_n$, for all x in \mathbb{U}_0 .

In the next step, we determine $\lim_{n\to\infty} \delta_n(\overline{x})$ and characterize the union of the sets $[x]_n$, $n \geq 0$.

Definition 2.5. For $x, y \in \mathbb{U}_0$, we define the relation " \sim_{∞} " on \mathbb{U}_0 as

$$x \sim_{\infty} y \Leftrightarrow \exists n \in \mathbb{N}_0 : T^n x = T^n y.$$

Further, put $[x]_{\infty} = \{z \in \mathbb{U}_0 : \exists n \in \mathbb{N}_0 \text{ such that } T^n z = T^n x\}$, and $\delta_{\infty}(x) = \min[x]_{\infty}$.

Lemma 2.12. The following holds.

1. The relation ' \sim_{∞} ' is an equivalence relation on \mathbb{U}_0 and, for all $x \in \mathbb{U}_0$, the set $[x]_{\infty}$ is the equivalence class of x with respect to this equivalence relation. Further, $[x]_{\infty} = [\delta_{\infty}(x)]_{\infty}$, and

$$[x]_{\infty} = \bigcup_{n>0} [x]_n.$$

2. For all $x \in \mathbb{U}_0$,

$$\delta_{\infty}(x) = \lim_{n \to \infty} \delta_n(x).$$

Proof. Ad 1. This is easily verified.

Ad 2. Let $\delta'(x) = \lim_{n \to \infty} \delta_n(x)$. Due to the fact that we are dealing with a convergent integer sequence, there exists an integer N such that for all $n \geq N$, $\delta'(x) = \delta_n(x)$. From the fact $\delta_n(x) \in [x]_n$, it follows that $\delta'(x) \in [x]_\infty$. Hence, $\delta_\infty(x) \leq \delta'(x)$.

On the other hand, for any $z \in [x]_{\infty}$, there exists $n \in \mathbb{N}_0$ such that $z \in [x]_n$. This implies that $z \geq \delta_n(x) \geq \delta'(x)$. We note that, by definition, $\delta_{\infty}(x) \in [x]_{\infty}$. Consequently, $\delta_{\infty}(x) \geq \delta'(x)$.

Corollary 2.13. The sets $[x]_{\infty}$, $x \in \mathbb{U}_0$, form a partition of \mathbb{U}_0 , $\mathbb{U}_0 = \bigcup_{x \in \mathbb{U}_0} [x]_{\infty}$. The 3x + 1 conjecture is equivalent to each of the following statements:

- 1. $\mathbb{U}_0 = [1]_{\infty}$.
- 2. $\forall x \in \mathbb{U}_0: \quad \delta_{\infty}(x) = 1.$

Let us study the action of T on the sets $[x]_{\infty}$.

Lemma 2.14. For all $x \in \mathbb{U}_0$,

$$T^{-1}[Tx]_{\infty} = [x]_{\infty}.$$

Proof. We have the following chain of equivalences:

$$z \in T^{-1}[Tx]_{\infty} \Leftrightarrow Tz \in [Tx]_{\infty} \Leftrightarrow \exists n \ge 0 : Tz \in [Tx]_n$$

$$\Leftrightarrow \exists n \ge 0 : T^{n+1}z = T^{n+1}x$$

$$\Leftrightarrow \exists n \ge 0 : z \in [x]_{n+1} \Leftrightarrow z \in [x]_{\infty}.$$

Corollary 2.15. In analogy to Corollary 2.9, for all $x \in \mathbb{U}_0$, we have

$$T[x]_{\infty} = [Tx]_{\infty}, \quad T^{-1}[x]_{\infty} = [\tau(x)]_{\infty}, \quad T[\tau(x)]_{\infty} = [x]_{\infty}.$$

Remark 2.5. For the analog equivalence classes on the set \mathbb{U} instead of \mathbb{U}_0 , one has $T[x]_{\infty} \subset [Tx]_{\infty}$, i.e., strict inclusion, as the inverse image $T^{-1}\{z\}$ of a point $z \in [Tx]_{\infty}$ may be empty.

In view of Lemma 2.14 and Corollary 2.15 we may introduce the following map.

Definition 2.6. Let $\mathbb{U}_{0,\infty}$ denote the quotient space $\mathbb{U}_0/\sim_{\infty}$, i.e., the set of equivalence classes associated with the equivalence relation ' \sim_{∞} '. The induced map T^* on $\mathbb{U}_{0,\infty}$ is defined as

$$T^*[x]_{\infty} = [Tx]_{\infty}, \quad x \in \mathbb{U}_0.$$

Theorem 2.16. The map T^* on $\mathbb{U}_{0,\infty}$ has the following properties:

- 1. T^* is well defined.
- 2. T^* is a bijection on $\mathbb{U}_{0,\infty}$.
- 3. T^* has the unique fixed point $[1]_{\infty}$:

$$[x]_{\infty} = T^*[x]_{\infty} \Rightarrow [x]_{\infty} = [1]_{\infty}.$$

4. It x is a periodic point of T, $T^k x = x$, with $k \ge 1$ the minimum period of x, then $[x]_{\infty}$ is a periodic point of $\mathbb{U}_{0,\infty}$ with $T^{*k}[x]_{\infty} = [x]_{\infty}$.

Proof. Ad 1. We have to show that the value of T^* is independent of the representative of the equivalence class $[x]_{\infty}$. Suppose that $[x]_{\infty} = [z]_{\infty}$. Then there exists $n \geq 0$ such that $T^n x = T^n z$. Hence, $T^{n+1} x = T^{n+1} z$, which implies $Tz \in [Tx]_n \subset [Tx]_{\infty}$. Thus, $[Tx]_{\infty} = [Tz]_{\infty}$.

Ad 2. By Lemma 2.14, $T^*[x]_{\infty} = T^*[z]_{\infty}$ implies $[Tx]_{\infty} = [Tz]_{\infty}$. Hence, there exists $n \geq 0$ such that $T^n(Tx) = T^n(Tz)$. As a consequence, $z \in [x]_{n+1} \subset [x]_{\infty}$, which implies $[z]_{\infty} = [x]_{\infty}$. Thus, T^* is injective on $\mathbb{U}_{0,\infty}$.

Let $[y]_{\infty}$ be an arbitrary element of $\mathbb{U}_{0,\infty}$. Then $[y]_{\infty} = [T\tau(y)]_{\infty} = T^*[\tau(y)]_{\infty}$. Thus, T^* is surjective.

Ad 3. Suppose that $[x]_{\infty} = T^*[x]_{\infty} = [Tx]_{\infty}$. Then there exists $n \geq 0$ such that $T^n x = T^n(Tx) = T(T^n x)$. As a consequence, the element $T^n x$ is a fixed point in \mathbb{U}_0 under T. This implies $T^n x = 1$ (see Remark 2.3, Part 3.). Due to $T^n 1 = 1$, we have $[x]_{\infty} = [1]_{\infty}$.

Ad 4. The property $T^k x = x$ implies $[T^k x]_{\infty} = T^{*k}[x]_{\infty} = [x]_{\infty}$.

3 Appendix

The idea underlying our approach to the 3x + 1 conjecture was to find a suitable metric space X in the form of some quotient space $X = \{\overline{x} : x \in \mathbb{U}_0\}$, and a contraction T^* on X that is intrinsically related to the map T in the sense that convergence of the sequences $(T^{*k}\overline{x})_{k\geq 0}$, $\overline{x} \in X$, to the unique fixed point of T^* implies the convergence of the sequences $(T^kx)_{k\geq 0}$ to 1, i.e., the validity of the 3x + 1 conjecture.

We were unable to realize this 'dream' of applying the Banach fixedpoint theorem, because we have not found an appropriate pair (X, T^*) . For example, $X = \mathbb{U}_{0,\infty}$ can easily be made into a metric space but it is the proof of the contraction property of the induced map T^* with respect to the chosen metric where we failed.

The following concepts allow a somewhat deeper understanding of the dynamics of the map T.

Definition 3.1. The *invertible* accelerated Collatz function $f: \mathbb{U}_0 \to \mathbb{U}_0$ is defined as follows. For $x \in \mathbb{U}_0$ and for $k \geq 0$, define $f^k x = T^k x$. For k < 0, put $f^k x = \tau^{-k}(x)$.

Consider the following equivalence relation on \mathbb{U}_0 :

$$x \sim y \Leftrightarrow \exists m, n \ge 0 : T^m x = T^n y.$$

We write [x] for the equivalence class of $x \in \mathbb{U}_0$, and get the following.

Lemma 3.1. Let the relation ' \sim' be defined as above and write \mathbb{U}'_0 for the quotient space \mathbb{U}_0/\sim . Then

1. For all $x \in \mathbb{U}_0$, the sets [x] are T-invariant in the following sense:

$$T^{-1}[x] = [x], \quad T[x] = [x] = [Tx].$$

2. For all $x \in \mathbb{U}_0$,

$$[x] = \bigcup_{k \in \mathbb{Z}} [f^k x]_{\infty} = \dots \cup [\tau(x)]_{\infty} \cup [x] \cup [Tx] \cup \dots$$

- 3. We have $[1] = [1]_{\infty}$.
- 4. The map $T': \mathbb{U}'_0 \to \mathbb{U}'_0$, T'[x] = [Tx] is well-defined and every element [x] of \mathbb{U}'_0 is a fixed point of T'.

Proof. The proof is straightforward and employs the techniques introduced in Section 2. \Box

Remark 3.1. The reader should note the behavior of the class [1], which is remarkably different from all other classes.

Remark 3.2. For $x \in \mathbb{U}_0$, put $\delta(x) = \min[x]$. The 3x + 1 conjecture is equivalent $\mathbb{U}_0 = [1]$. Further, it is equivalent to $\delta(x) = 1$ for all $x \in \mathbb{U}_0$.

Remark 3.3. Lemma 3.1 tells us that every set [x] is T-invariant, which is to say that $T^{-1}[x] = [x]$. In addition to this result, Theorem 2.16 shows that $[1] = [1]_{\infty}$ is the only T-invariant set of the form $[x]_{\infty}$. These two results call out for the application of concepts from the theory of dynamical systems, for example from ergodic theory. Let (X, \mathcal{B}, m) be a probability space. A measure preserving map $f: X \to X$ is called ergodic if the only f-invariant elements A of \mathcal{B} , i.e., $f^{-1}A = A$, are those with m(A) = 0 or m(A) = 1. It is well known that ergodicity of f is equivalent to each of the following properties: (i) for every $A \in \mathcal{B}$ with m(A) > 0 we have $m(\bigcup_{k=1}^{\infty} f^{-k}A) = 1$, or (ii) for every $A, B \in \mathcal{B}$ with m(A) > 0 and m(B) > 0, there exists k > 0 with $m(f^{-k}A \cap B) > 0$ (see, for example, Walters[4, Theorem 1.5]). In our case, we would have to prove ergodicity for f = T, where \mathbb{U}_0 would have to be equipped with an appropriate probability space structure. We would then be able to derive the 3x + 1 conjecture for almost all x.

The following two notions allow some kind of "bookkeeping" when we iterate the map f. With every $x \in \mathbb{U}_0$, we may associate two infinite matrices as follows.

Definition 3.2. Let $x \in \mathbb{U}_0$. We define the matrix of equivalence classes associated with x as $C(x) = (c_{k,n})_{k \in \mathbb{Z}, n \geq 0}$, where $c_{k,n} = [f^k x]_n$. In addition, we define the matrix of minimal elements associated with x

In addition, we define the matrix of minimal elements associated with x as $M(x) = (\mu_{k,n})_{k \in \mathbb{Z}, n \geq 0}$, with $\mu_{k,n} = \min c_{k,n}$. Let $\delta^*(x)$ denote the minimal element of the matrix M(x).

Clearly, we have $\mu_{k,n} = \delta_n(f^k x)$, and $\delta^*(x) = \delta(x)$. Note that if we fix the row index k, then the row $(\mu_{k,n})_{n\geq 0}$ in M(x) has a constant tail eventually, because the convergent sequence $(\delta_n(f^k x))_{n\geq 0}$ is constant from some index N = N(k) onwards, with every element then being equal to $\delta_{\infty}(f^k x)$.

There is even further 'tail'-structure in M(x): suppose that $\delta^*(x)$ is equal to $\mu_{k,n}$, where k and n are minimal with this property (in this order). Then

 $\delta^*(x) = \mu_{k,m}$ for all $m \ge n$. That is to say, the k-th row becomes eventually constant.

From the discussion above it follows that it is sufficient to prove the 3x + 1 conjecture for the subset $\{\delta_{\infty}(x) : x \in \mathbb{U}_0\}$ of \mathbb{U}_0 or, alternatively, $\{\delta_1(x) : x \in \mathbb{U}_0\}$. These facts suggest the following notion.

Definition 3.3. A subset V of \mathbb{U}_0 is called sufficient if the validity of the 3x + 1 conjecture for every element of V implies the validity of the 3x + 1 conjecture for every element of \mathbb{U}_0 .

Lemma 3.2. The set $\{x \in \mathbb{U}_0 : 1 \leq \nu_2(3x+1) \leq 4\}$ is sufficient.

Proof. Let $x \in \mathbb{U}_0$ be arbitrary. Trivially, we have $[x]_{\infty} = [\delta_1(x)]_{\infty}$. As a consequence, the set $\{\delta_1(x) : x \in \mathbb{U}_0\}$ is sufficient. Further, $\delta_1(x) = \tau(Tx)$.

From Lemma 2.3 it follows that $\nu_2(3\xi(x)+1) \in \{1,2\}$, for all $x \in \mathbb{U}_0$. Due to the fact that either $\tau(x) = \xi(x)$, or $\tau(x) = S\xi(x)$, we have $\nu_2(3\tau(x)+1) \in \{1,2,3,4\}$.

The reader might want to compare this result with Sander [3, Theorem 1]. For further, very extensive results on sufficient sets we refer the reader to Monks[2].

Acknowledgements

The author would like to thank Harry (Hillel) Furstenberg, who brought this problem to his attention in several personal discussions some decades ago at CIRM in Luminy, France.

References

- [1] Jeffrey C. Lagarias, editor. The Ultimate Challenge: The 3x+1 Problem. American Mathematical Society, Providence, RI, 2010.
- [2] Keenan Monks, Kenneth G. Monks, Kenneth M. Monks, and Maria Monks. Strongly sufficient sets and the distribution of arithmetic sequences in the 3x + 1 graph. Discrete Math., 313(4):468-489, 2013.
- [3] J. W. Sander. On the (3N + 1)-conjecture. Acta Arith., 55(3):241-248, 1990.
- [4] Peter Walters. An Introduction to Ergodic Theory, volume 79 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1982.

Author's address:

Peter Hellekalek, Dept. of Mathematics, University of Salzburg, Hellbrunner-strasse 34, 5020 Salzburg, Austria

E-mail: peter.hellekalek@sbg.ac.at