



Updates to the Evidence to Recommendation Framework: Pfizer-BioNTech and Moderna COVID-19 vaccine booster doses

Sara Oliver, MD, MSPH
ACIP Meeting
November 19, 2021



cdc.gov/coronavirus

Evidence to Recommendations (EtR) Framework



Evidence to Recommendations (EtR) Framework

- Previous presentations/discussions for booster doses of COVID-19 vaccines

September 23rd:

COVID-19 vaccine booster doses: Benefit/risk discussion

Evidence to Recommendation Framework: Booster doses of Pfizer-BioNTech COVID-19 vaccine

VOTE: Pfizer-BioNTech COVID-19 booster doses

<https://www.cdc.gov/vaccines/acip/meetings/slides-2021-09-22-23.html>

October 21st:

National Institutes of Health: Mix and Match booster study

Evidence to Recommendation Framework: Booster doses of Moderna & Janssen COVID-19 vaccines

VOTE: Moderna & Janssen COVID-19 booster doses (including heterologous boosting)

<https://www.cdc.gov/vaccines/acip/meetings/slides-2021-10-20-21.html>

COVID-19 vaccine booster dose in persons who received a Janssen COVID-19 vaccine primary dose

- Persons aged ≥ 18 years who received primary vaccination with Janssen COVID-19 vaccine **should** receive a single COVID-19 vaccine booster dose at least 2 months later
- Any FDA-approved or authorized COVID-19 vaccine (Pfizer-BioNTech, Moderna, or Janssen) can be used for booster dose, regardless of vaccine received for primary series

COVID-19 vaccine booster dose in persons who completed an mRNA primary series

Persons who should receive a COVID-19 booster dose

- Aged ≥ 65 years
- Aged ≥ 18 years and reside in long-term care settings
- Aged 50-64 years with certain underlying medical conditions

Persons who may receive a COVID-19 booster dose, based on individual benefits and risks

- Aged 18-49 years with certain underlying medical conditions*
- Aged 18-64 years at increased risk for SARS-CoV-2 exposure and transmission because of occupational or institutional setting

- Booster dose administered at least 6 months after completion of primary series
- Any FDA-approved or authorized COVID-19 vaccine (Pfizer-BioNTech, Moderna, or Janssen) can be used for booster dose, regardless of vaccine received for primary series

* Includes pregnant people

COVID-19 vaccine booster dose in persons who completed an mRNA primary series

Persons who should receive a COVID-19 booster dose

- Aged ≥ 65 years
- Aged ≥ 18 years and reside in long-term care settings
- Aged 50-64 years with certain underlying medical conditions

Persons who may receive a COVID-19 booster dose, based on individual benefits and risks

- Aged 18-49 years with certain underlying medical conditions*
- Aged 18-64 years at increased risk for SARS-CoV-2 exposure and transmission because of occupational or institutional setting

- Booster dose administered at least 6 months after completion of primary series
- Any FDA-approved or authorized COVID-19 vaccine (Pfizer-BioNTech, Moderna, or Janssen) can be used for booster dose, regardless of vaccine received for primary series

* Includes pregnant people

Policy Question

- Do the balance of benefits and risks and facilitation of implementation warrant an update to COVID-19 vaccine policy?


All other persons ≥ 18 years of age **may** receive a COVID-19 booster dose ≥ 6 months after completion of the mRNA primary series under the current Emergency Use Authorization

Persons who may receive a COVID-19 booster dose, based on individual benefits and risks

- Aged 18-49 years with certain underlying medical conditions*
- Aged 18-64 years at increased risk for SARS-CoV-2 exposure and transmission because of occupational or institutional setting
- All other persons aged ≥ 18 years


Evidence to Recommendations Framework

Booster doses of COVID-19 vaccines




Trends in COVID-19 cases in the United States

January 23, 2020 – November 16, 2021




Trends in COVID-19 hospitalizations in the United States

August 1, 2020 – November 15, 2021



US COVID-19 7-day case rate per 100,000

By state/territory



7-Day Case Rate per 100,000

Data not available ●

0 - 48.9 ●

59.4 - 104.1 ●


118.3 - 159.3 ●

179.5 - 268.4 ●

287.3 - 371.9 ●

402.6 - 589.3 ●

Number of people fully vaccinated in the U.S. by COVID-19 vaccine series type




Over **195** million people fully vaccinated in the US

69% of the population ≥ 12 years of age


COVID-19 vaccine first doses administered, by age group

December 14, 2020 – November 15, 2021



Source: Immunization Data Lake.

VE against symptomatic infection by age group and time since vaccination in pre-Delta and Delta periods




— Pre-Delta (March 13–May 29) with 95% CIs in dotted lines
— Delta (July 18–August 31) with 95% CIs in dotted lines

- VE is lower during Delta
- VE wanes during both periods
- Curves similar across age groups
- For ≥ 65 , VE lower than for other age groups soon after vaccination, no clear trend over time since vaccination

The presented (fitted) curves are truncated on the day with ≤ 10 cases observed beyond it to avoid presenting wide confidence bounds.


Vaccine effectiveness for Pfizer-BioNTech vaccine by time since second dose, outcome, and age

Author, study platform, variant(s), outcome, age




Vaccine effectiveness for Pfizer-BioNTech vaccine by time since second dose, outcome, and age

Author, study platform, variant(s), outcome, age



Vaccine effectiveness for Pfizer-BioNTech vaccine by time since second dose, outcome, and age

Author, study platform, variant(s), outcome, age




Documented or symptomatic infection

Severe disease, hospitalization, death



Vaccine effectiveness for Moderna vaccine by time since second dose, outcome, and age

Author, study platform, variant(s), outcome, age



Summary

- Over 195 million people are fully vaccinated in the United States
- COVID-19 cases are increasing in some jurisdictions recently
- VE after primary series waning for infection, but protection remains high for severe disease and hospitalization
 - Waning appears to be less pronounced for Moderna COVID-19 vaccine, compared to Pfizer-BioNTech COVID-19 vaccine recipients

Evidence to Recommendations Framework

Booster doses of COVID-19 vaccines



PICO Question

	Pfizer-BioNTech	Moderna
Population	Persons aged ≥ 18 years who completed a COVID-19 vaccine primary series ≥ 6 months ago	
Intervention	Pfizer-BioNTech COVID-19 Vaccine booster dose (BNT162b2, 30 μg , IM)	Moderna COVID-19 Vaccine booster dose (mRNA-1273, 50 μg , IM)
Comparison	No booster dose	
Outcomes	Symptomatic laboratory-confirmed COVID-19* Hospitalization due to COVID-19* Death due to COVID-19 Transmission of SARS-CoV-2 infection Serious adverse events* Reactogenicity	

* Critical outcomes

Updated Pfizer-BioNTech booster data

- Phase 3 booster dose randomized control trial (RCT)
- ~10,000 participants from phase 2/3 efficacy trial
 - All participants received 2-dose primary series of Pfizer-BioNTech COVID-19 vaccine
- Pfizer-BioNTech booster dose and placebo doses randomized 1:1
 - Randomization was stratified by age
 - 60% 16-55 years
 - 40% >55 years
 - Booster doses given 10-12 months after second dose
- Median follow-up of 2.5 months post booster dose

Outcome 1: Symptomatic laboratory-confirmed SARS-CoV-2 infection

Population	Events/Vaccine (n/N)	Events/Placebo (n/N)	Relative vaccine efficacy (95% CI)
Primary Outcome			
No evidence of prior infection, ≥7 days post booster	6/4659	123/4614	95.2% (89.3%, 97.9%)
Secondary Outcomes			
± evidence of prior infection, ≥7 days post booster	7/4994	124/4963	94.5% (88.3%, 97.4%)
All available efficacy (± evidence of prior infection, post booster)	15/5003	141/4943	89.8% (82.6%, 94.0%)
0-7 days post booster	8/5003	15/4943	47.4% (-24.0%, 77.7%)
≥7 days to <2 months post booster	6/4995	112/4928	94.9% (88.5%, 97.8%)
≥2 months to <4 months post booster	1/4891	14/4616	93.3% (48.9%, 99.1%)

Outcome 1: Symptomatic laboratory-confirmed SARS-CoV-2 infection

Population	Events/Vaccine (n/N)	Events/Placebo (n/N)	Relative vaccine efficacy (95% CI)
Primary Outcome			
No evidence of prior infection, ≥7 days post booster	6/4659	123/4614	95.2% (89.3%, 97.9%)
Secondary Outcomes			
± evidence of prior infection, ≥7 days post booster	7/4994	124/4963	94.5% (88.3%, 97.4%)
All available efficacy (± evidence of prior infection, post booster)	15/5003	141/4943	89.8% (82.6%, 94.0%)
0-7 days post booster	8/5003	15/4943	47.4% (-24.0%, 77.7%)
≥7 days to <2 months post booster	6/4995	112/4928	94.9% (88.5%, 97.8%)
≥2 months to <4 months post booster	1/4891	14/4616	93.3% (48.9%, 99.1%)

Evaluation of other beneficial outcomes

- Hospitalization due to COVID-19
 - No hospitalizations due to COVID-19 occurred in the booster (3-dose) or placebo (2-dose) groups
- Death due to COVID-19
 - No deaths due to COVID-19 occurred in the booster (3-dose) or placebo (2-dose) groups
- Transmission of SARS-CoV-2 infection
 - No data to assess this outcome

Outcome 6: Serious Adverse Events

Study/population ^a	Events/Vaccine (n/N)	% SAE Vaccine	Events/Placebo (n/N)	% SAE Placebo	Associated with vaccination ^b
Pfizer/BioNTech, unpublished	16/5055	0.3	24/5020 ^c	0.5	3

a. Included all randomized participants who received a booster dose

b. Three serious adverse events among booster recipients were deemed by blinded investigators to be related to vaccination. These included: moderate persistent tachycardia, moderate transient elevated hepatic enzymes, and mild elevated hepatic enzymes.

c. There was one death among placebo recipients and none among booster dose recipients

Adverse events of clinical interest

- No cases of anaphylaxis, hypersensitivity, or myocarditis/pericarditis reported
 - Given the rarity of these adverse events, we would not expect to capture them in a RCT of this size
- Lymphadenopathy was more common after the 3rd dose (**2.7%**) than after the primary series (**0.4%**)
 - Typically mild to moderate and located in the axilla or cervical nodes
 - Most occurred 1-3 days post booster and resolved within 1-3 days of onset
 - Frequency higher in younger participants and female participants

Outcome 7: Reactogenicity

- No updated data from phase 3 booster trial

Pfizer-BioNTech booster ≥ 6 months after primary series

Changes to GRADE from previous booster discussion

Outcome	Importance	Previous evidence certainty	Current evidence certainty
Benefits			
Symptomatic laboratory-confirmed COVID-19	Critical	Very low	High
Hospitalization due to COVID-19	Critical	Very low	Very low
Death due to COVID-19	Important	No data	No data
Transmission of SARS-CoV-2 infection	Important	No data	No data
Harms			
Serious adverse events	Critical	Very low	Low
Reactogenicity	Important	Very low	Very Low

Myocarditis in Israel

Reported after Pfizer-BioNTech COVID-19 vaccine, December 2020-October 10, 2021

	Age (years)	Post-dose 1 Rate per 100,000	Post-dose 2 Rate per 100,000	Post-dose 3 Rate per 100,000	Number of 3 rd dose delivered
Females	12-15	0	0.6	0	279
	16-19	0	0.9	0	97,807
	20-24	0.4	2.5	0	141,910
	25-29	0	0.4	0	130,283
	≥30	0.1	0.3	0	1,542,142
Males	12-15	0.5	6.6	0	292
	16-19	1.2	16.1	5.2	96,238
	20-24	2.2	10.3	3.6	139,015
	25-29	1.2	8.4	0.7	133,650
	≥30	0.5	1.7	0.4	1,448,745

Rates of myocarditis after a third dose appear to fall **between** rates seen after **dose 1** and **dose 2**


Benefits and risks after Pfizer-BioNTech COVID-19 booster dose

For every million doses of vaccine given

Scenario:

- VE for hospitalization averaged from four platforms¹
- Boost to 95% VE for hospitalization
- **Myocarditis risk equivalent to risk after 1st and 2nd dose averaged**

Age Group	VE for hospitalization
18 – 29 years	90.7%
30 – 49 years	90.2%
50 – 64 years	91.1%
≥65 years	85.1%



1. Scobie et al., COVID-NET, VISION, IVY Network

COVID-NET hospitalization rates from the week of August 21, 2021; Myocarditis rates from VAERS data through August 18, 2021

Benefits and risks after Pfizer-BioNTech COVID-19 booster dose

- Phase 3 RCT booster efficacy data demonstrates booster dose provides additional protection and is safe
 - No hospitalized cases of COVID-19 after 2-dose primary series
- Based on data from Israel, myocarditis risk after booster dose of Pfizer-BioNTech COVID-19 vaccine appears to fall between rates seen after dose 1 and dose 2

Moderna booster data

No booster vaccine efficacy/effectiveness studies identified

Moderna booster study previously presented*

Part A: Phase 2 randomized, observer-blind, placebo-controlled dose confirmation study among participants aged ≥ 18 years

Part B: study amended for open label phase based on participant selection

Placebo recipients \rightarrow 100 μg primary series

50 μg primary series \rightarrow 50 μg booster (≥ 6 months after dose 2)

100 μg primary series \rightarrow 50 μg booster (≥ 6 months after dose 2)

Immunobridging to patients in Phase 3 efficacy study

Prespecified non-inferiority analysis

28 days after booster versus 28 days after dose 2 of primary series

Moderna booster data

- Symptomatic laboratory-confirmed COVID-19
 - Geometric mean titers of booster recipients (N=149) compared to primary series recipients (N=1,053)
 - The geometric mean ratio of 1.76 (95%CI: 1.50–2.06) met non-inferiority criteria
- Serious Adverse Events
 - No SAEs occurred among the 171 participants receiving a 50 ug booster in the open label booster study within 28 days of the booster dose
- Reactogenicity
 - Severe reactogenicity occurred among 10.8% of participants receiving a booster in the open label booster study

Moderna booster ≥ 6 months after primary series


Summary of GRADE – No update to previous discussion

Outcome	Importance	Evidence certainty
Benefits		
Symptomatic laboratory-confirmed COVID-19	Critical	Very low
Hospitalization due to COVID-19	Critical	No data
Death due to COVID-19	Important	No data
Transmission of SARS-CoV-2 infection	Important	No data
Harms		
Serious adverse events	Critical	Very low
Reactogenicity	Important	Very low

Benefits and risks after Moderna COVID-19 booster dose


- No phase 3 RCT booster efficacy data available for Moderna
 - Immunogenicity study demonstrates the ability to boost antibody levels
- Effectiveness after a primary series appears to have waned less in Moderna than in Pfizer
- Myocarditis risk after booster dose of Moderna is unknown
 - Accumulating evidence from multiple sources suggests a higher risk for myocarditis following Moderna compared to Pfizer-BioNTech primary series vaccination
 - Moderna booster dose is a lower dose (50µg) than the primary series dose (100µg)

Number needed to vaccinate with booster dose to prevent one hospitalization over 6 months



** Not estimable due to pre-booster efficacy estimated at >95%

Number needed to vaccinate with booster dose to prevent one infection over 6 months



Summary of safety surveillance findings

- V-safe
 - For Pfizer-BioNTech and Moderna, local and systemic reactions were reported less frequently following a booster dose than dose 2 of the primary series
 - Moderna booster appears to be more reactogenic than Pfizer-BioNTech booster, regardless of the primary series manufacturer
- VAERS
 - Most reports ($\geq 93\%$) were non-serious (similar to primary series)
 - Most frequently reported non-serious AEs were known and well characterized AEs associated with COVID-19 vaccination
 - 54 preliminary reports of myocarditis
 - 12 verified reports that met CDC case definition

Impact of a booster dose on transmission

- After a primary mRNA COVID-19 vaccine series, protection against **asymptomatic infection** (and presumably transmission) was found for a time period^{1,2}
 - Largest impact seen in the first 2 months post-vaccination²
 - Likely an impact of very high antibody titers
- Limited data on impact of booster dose on asymptomatic infection/transmission
 - One study from Israel found **lower viral loads** in patients with breakthrough infections after a booster dose, similar to viral loads seen within 2 months after primary series²
 - Early VE against SARS-CoV-2 infection after a booster dose demonstrates increase in VE (including asymptomatic infection)^{3,4}
- While protection against asymptomatic infection may not be permanent, even temporary protection may favor into benefit/risk balance approaching winter and holidays with increased travel and indoor gatherings

¹CDC Science Brief <https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/fully-vaccinated-people.html> ²Levine-Tiefenbrun et al. Nature Medicine 2021 <https://www.nature.com/articles/s41591-021-01575-4>

³Saciuk et al. Journal of Infectious Diseases <https://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiab556/6415586>

⁴Andrews et al. [Effectiveness of BNT162b2 \(Comirnaty, Pfizer-BioNTech\) COVID-19 booster vaccine against covid-19 related symptoms in England: test negative case-control study | medRxiv](https://doi.org/10.1101/2021.09.14.21260000)

Summary

Balance of benefits and harms for booster doses

- Booster dose of Pfizer-BioNTech COVID-19 vaccine is **effective** in preventing laboratory confirmed symptomatic SARS-CoV-2
- Data from Moderna trial does not provide efficacy data, but demonstrates the ability to boost immune response
- Individual benefit/risk balance for booster doses of an mRNA vaccine **varies by age**
 - Older adults have the clearest benefit/risk balance
 - Among other ages, variation within balance of benefits and risks
 - Myocarditis data after booster doses reassuring to date
- Unable to account for other benefits
 - Possible impact on rates of community transmission


Evidence to Recommendations Framework

Booster doses of COVID-19 vaccines




Survey of fully vaccinated adults (vaccine booster eligibility)

- Survey respondents were asked:
 - “Have you personally received a booster or additional dose of the COVID-19 vaccine after you were already fully vaccinated?”
 - **4 in 10** fully vaccinated adults are **unsure** whether they are eligible for a booster dose




Vaccine booster confidence

- Survey respondents were asked:
 - “Which comes closer to your view about the news that some people might need vaccine boosters?”
 - **More than 6 in 10** adults overall say the news that some people might need boosters “**shows that scientists are continuing to find ways to make vaccines more effective.**”





Evidence to Recommendations Framework

Booster doses of COVID-19 vaccines



Number of people with a booster dose in the U.S. by COVID-19 vaccine series type

Over **31** million people have received a booster dose in the US




Daily number of booster doses administered, by manufacturer

Persons ≥18 years of age

Daily Additional Doses (K)

Cumulative Additional Doses Administered (M)



Note: Number of people who received a booster dose includes anyone who is fully vaccinated and has received another dose of COVID-19 vaccine starting August 13, 2021. This includes people who received booster doses and people who received additional doses.

Source: Immunization Data Lake. **Data as of November 16, 2021, 0600AM.**

Implementation of COVID-19 vaccine booster doses

- At least **31 million** individuals in the United States have received a COVID-19 vaccine booster dose
- ~**17 million** individuals ≥ 65 years of age received a COVID-19 booster dose
- Some states currently broadening booster eligibility criteria


Implementation of COVID-19 vaccine booster doses

Considerations


- Vaccine recommendations based on risk and exposures are more difficult to implement than age-based recommendations
- ACIP recommendations that are **consistent** with the FDA EUA are easier to communicate and implement
- If recommendations for booster doses varied across the two mRNA vaccines, it would be difficult to communicate and implement

Evidence to Recommendations Framework

Booster doses of COVID-19 vaccines




Percentage of people who have received at least one dose of the COVID-19 vaccine by race/ethnicity over time



Booster doses in persons ≥ 18 years of age with completed vaccination series ≥ 6 months earlier* by race/ethnicity

Data from 27,085,156 people with booster doses
 Race/ethnicity was available for 18,832,653 (69.5%) people with a booster dose



Notes: *Includes Janssen booster doses for people who have completed a primary vaccination series ≥ 2 months earlier. Number of people who received a booster dose includes anyone who is fully vaccinated and has received another dose of COVID-19 vaccine starting August 13, 2021. This includes people who received booster doses and people who received additional doses. The expected timing for a booster dose was set at ≥ 6 months after primary series completion. Primary series is determined by the vaccine type of the second mRNA dose received or the first J&J/Janssen dose received. **Does not include vaccine administrations reported by Texas as the primary series cannot be linked to booster dose in the aggregate format submitted by Texas.** Source: Immunization Data Lake. Data as of November 16, 2021, 0600AM.

Summary - Equity

- Some disparities in primary series vaccine delivery have improved over time
- Early data on COVID-19 booster doses demonstrates disparities by race and ethnicity
 - Recommendations that are complex, difficult to communicate, or difficult to implement may worsen disparities in booster vaccination rates

Summary



Work Group Interpretation

- **Top priority** should be **continued vaccination of unvaccinated individuals**
- Balance of benefits and risks **varies by age**
 - Older adults have the clearest benefit/risk balance
 - Myocarditis data after booster doses reassuring to date, continue to closely monitor
 - Increases in COVID-19 cases may also impact benefit/risk balance

Goals of COVID-19 vaccines:

- Primary goal: Prevention of **severe disease**
- Secondary goals:
 - Maintaining workforce and healthcare capacity
 - Reduce infection and transmission
- Unknown impact of COVID-19 vaccine booster dose on prevention of transmission.
However, even reduction in transmission may be important around winter and holidays

Evidence to Recommendations Framework

Type of recommendation	We do not recommend the intervention	We recommend the intervention for individuals based on assessment of benefits and risks	We recommend the intervention
-------------------------------	--------------------------------------	---	-------------------------------



Used when the **risks** clearly outweigh the **benefits**

Used when there is diversity of the **benefits** and **risks**

Can allow flexibility across a population

Used when the **benefits** clearly outweigh the **risks**

Evidence to Recommendations Framework

Type of recommendation	We do not recommend the intervention	We recommend the intervention for individuals based on assessment of benefits and risks	We recommend the intervention
-------------------------------	--------------------------------------	---	-------------------------------



Used when the **risks** clearly outweigh the **benefits**

Used when there is diversity of the **benefits** and **risks**

Used when the **benefits** clearly outweigh the **risks**

Can allow flexibility across a population

MAY
receive a booster

SHOULD
receive a booster

Individual benefit-risk considerations for people who may receive a booster dose

- Potential **benefits** of booster dose
 - Reduced risk of SARS-CoV-2 infection, severe disease
 - May reduce transmission of SARS-CoV-2 to others
- Potential **risks** of booster dose
 - Rare risks of serious adverse events (e.g., myocarditis, pericarditis, TTS, GBS, anaphylaxis)
 - Common risks of transient local and systemic symptoms
- **Individual risk factors** for SARS-CoV-2 infection
 - Risk of exposure (occupational and institutional settings, e.g., healthcare workers, long term care settings)
 - Risk for infection (time since completion of primary series)
- **Individual impacts** of SARS-CoV-2 infection
 - Risk for severe infection (related to underlying conditions)
 - Risk associated with a person's circumstances (living with/caring for at-risk individuals or consequences of inability to meet obligations due to infection)

Updates for the future

- Updates from a Moderna COVID-19 booster dose
- Rates of myocarditis after 3rd dose
- Updated data for overall safety profile of booster doses
- Continued evaluations for vaccine effectiveness
 - Includes VE for primary series and booster doses



Current recommendations for booster doses of COVID-19 vaccines

Age	mRNA COVID-19 vaccine primary series				Janssen COVID-19 vaccine primary series
	No risk factors	Occupational or institutional exposures	Underlying medical conditions	Resident of LTCF	
≥65 years	Should receive a booster			Should receive a booster	Should receive a booster
50–64 years	Not eligible	May receive a booster	Should receive a booster		
18–49 years			May receive a booster		



Proposed recommendations for booster doses of COVID-19 vaccines

Age	mRNA COVID-19 vaccine primary series			Janssen COVID-19 vaccine primary series
	No risk factors	Underlying medical conditions	Resident of LTCF	
≥65 years	Should receive a booster	Should receive a booster	Should receive a booster	Should receive a booster
50–64 years	May receive a booster			
18–49 years		May receive a booster		

Policy Question

- Do the balance of benefits and risks and facilitation of implementation warrant an update to COVID-19 vaccine policy?

All other persons ≥ 18 years of age **may receive** a COVID-19 booster dose ≥ 6 months after completion of the mRNA primary series under the current Emergency Use Authorization

Persons who should receive a COVID-19 booster dose

- Aged ≥ 65 years
- Aged ≥ 18 years and reside in long-term care settings
- Aged 50-64 years with certain underlying medical conditions

Persons who may receive a COVID-19 booster dose, based on individual benefits and risks

- All other persons aged ≥ 18 years



Proposed recommendations for booster doses of COVID-19 vaccines

Option #2

Age	mRNA COVID-19 vaccine primary series		Janssen COVID-19 vaccine primary series
	No risk factors	Resident of LTCF	
≥65 years	Should receive a booster	Should receive a booster	Should receive a booster
50–64 years			
18–49 years	May receive a booster		

COVID-19 vaccine booster dose in persons who completed an mRNA primary series: **PROPOSED, Option #2**

Persons who should receive a COVID-19 booster dose

Aged ≥ 50 years

Aged ≥ 18 years residing in LTCF

Persons who may receive a COVID-19 booster dose, based on individual benefits and risks

All other persons aged ≥ 18 years

- Booster dose administered at least 6 months after completion of primary series
- Any FDA-approved or authorized COVID-19 vaccine (Pfizer-BioNTech, Moderna, or Janssen) can be used for booster dose, regardless of vaccine received for primary series

ACIP Vote

Interim Recommendation

A single COVID-19 vaccine booster dose is recommended for persons aged **≥18 years*** who received an mRNA COVID-19 vaccine primary series based on **individual benefit and risk**, at least 6 months after the primary series, under the FDA's Emergency Use Authorization


*Individuals ≥18 years of age not otherwise recommended to receive a COVID-19 vaccine

ACIP Vote
Interim Recommendation
Option #2

A single COVID-19 vaccine booster dose is recommended for persons aged **≥50 years** who received an mRNA COVID-19 vaccine, at least 6 months after the primary series, under the FDA's Emergency Use Authorization

Acknowledgments

- Monica Godfrey
- Megan Wallace
- Sarah Mbaeyi
- Rachel Gorowitz
- Jack Gersten
- Jefferson Jones
- Eddie Shanley
- Mary Chamberland
- Deblina Datta
- Stephen Hadler
- VTF ACIP WG Team
- ACIP COVID-19 Vaccines Work Group
- Vaccine Task Force
- Epi Task Force
- Respiratory Viruses Branch



For more information, contact CDC
1-800-CDC-INFO (232-4636)
TTY: 1-888-232-6348 www.cdc.gov

Thank you

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

